



UNIVERSITI PUTRA MALAYSIA

**INFLUENCE OF HEPARIN AND SEMINAL HEPARIN BINDING
PROTEINS ON REPRODUCTIVE PERFORMANCE OF SAHIWAL-
FRIESIAN BULLS AND RABBITS**

RAMAKRISHNAN A/L PATCHAIAPPAN

FPV 2003 11

**INFLUENCE OF HEPARIN AND SEMINAL HEPARIN BINDING
PROTEINS ON REPRODUCTIVE PERFORMANCE OF SAHIWAL-
FRIESIAN BULLS AND RABBITS**

By

RAMAKRISHNAN A/L PATCHAIAPPAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia in Fulfilment of the Requirements for
the Degree of Doctor of Philosophy**

December 2003



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Doctor of Philosophy

**INFLUENCE OF HEPARIN AND SEMINAL HEPARIN BINDING
PROTEINS ON REPRODUCTIVE PERFORMANCE OF SAHIWAL-
FRIESIAN BULLS AND RABBITS**

By

RAMAKRISHNAN A/L PATCHAIAPPAN

December 2003

Chairman: Prof. Dato' Dr. Sheikh Omar Abdul Rahman

Faculty: Faculty of Veterinary Medicine

A study was undertaken to characterise seminal heparin binding proteins (HBP) of Sahiwal-Friesian bulls and rabbits and to investigate the influence of HBP on the reproductive performance of Sahiwal-Friesian bulls and rabbits.

Using one dimensional sodium dodecyl sulfate-polyacrylamide (SDS-PAGE) gel electrophoresis, the major classes of HBP identified and isolated from Sahiwal-Friesian bulls and rabbits were with molecular weights of 15, 25, 30, 38 and 42, 15, 25, 38, 42 and 101 kDa, respectively. Bulls of varying fertility were found to have similar classes of HBP in their seminal fluids in the present study. This may show that the presence of the different classes of HBP in seminal fluid may not be an indicative of bull fertility.

Heparin and lyophilised HBP were found to have no effect on the percentage of spermatozoa motility in Sahiwal-Friesian bulls and rabbits. However, the addition of heparin or lyophilised HBP to epididymal spermatozoa and ejaculated spermatozoa has significantly increased the rate of acrosome reaction in both the species.

The present study also showed that rabbits inseminated with epididymal spermatozoa without the addition of HBP produced significantly higher mean littered rates and mean litter born per doe as compared to those animals inseminated with epididymal spermatozoa treated with different levels of HBP. These results showed that epididymal spermatozoa of rabbit could fertilise oocytes without prior exposure to HBP or seminal fluid whose HBP were believed to be necessary for capacitation of spermatozoa in cattle and rabbit.

The group of rabbits inseminated with epididymal spermatozoa treated with HBP at the rate of 4 mg/ml, recorded significantly lower mean litter born per doe as compared to those female rabbits inseminated with epididymal spermatozoa treated with HBP at the lower rates. These results may show that a higher concentration of lyophilized HBP causes more damage to the epididymal spermatozoa instead of further improving their fertilising ability.

Among the four groups of rabbits inseminated with epididymal spermatozoa (ES), epididymal spermatozoa with HBP (ESII), ejaculated spermatozoa (ESM) and ejaculated spermatozoa with HBP (ESMH), there was no significant difference found for the litter rate between the groups inseminated at 0 hour. However, for animals inseminated at 6 hours after induction of ovulation, the litter rate was significantly lower in the group inseminated with ES as compared to the other three groups which were inseminated with spermatozoa which had either natural HBP or added lyophilised HBP. In the present study, it was also found that none of the female rabbits inseminated at 12 and 24 hours after the vaginal stimulation littered.

It was also observed that epididymal spermatozoa samples added with lyophilised HBP bound significantly higher concentration of tritiated (^3H) heparin to their membrane as compared to samples that were devoid of HBP. The present study also showed that higher fertile bulls bound significantly higher concentrations of (^3H) heparin to their spermatozoa membrane than the lower fertile bulls.

From this study, it can be concluded that (i) epididymal spermatozoa of rabbits could fertilise oocytes without prior exposure to HBP or seminal fluid in the rabbit and (ii) Sahiwal-Friesian bulls could be selected based on their ability to bind (^3H) heparin to their spermatozoa.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**PENGARUH HEPARIN DAN PROTEIN PELEKAT HEPARIN KE ATAS
PESTASI PEMBIAKAN LEMBU JANTAN SAHIWAL-FRIESIAN DAN
ARNAB**

Oleh

RAMAKRISHNAN A/L PATCHAIAPPAN

Disember 2003

Pengerusi: Professor Dato' Sheikh Omar Abdul Rahman, Ph.D.

Fakulti : Fakulti Perubatan Veterinar

Suatu kajian pencirian protein pelekat heparin (HBP) dalam semen dari lembu jantan Sahiwal-Friesian dan arnab telah dijalankan. Pengaruh HBP ke atas perstasi pembiakan lembu jantan Sahiwal-Friesian dan arnab juga dikaji.

Dengan menggunakan dimensi satu sodium dodecyl sulfate-polyacrylamide (SDS-PAGE) gel electrophoresis, kelas utama HBP dalam semen telah dikenalpasti dan diasingkan dari lembu jantan Sahiwal-Friesian dan arnab masing-masing mempunyai berat molekul 15, 25, 30, 38 dan 42, 15, 25, 38, 42 dan 101 kDa. Dari kajian ini didapati bahawa lembu jantan yang mempunyai kesuburan yang berbeza mengandungi kelas HBP yang sama dalam semen. Ini mungkin menunjukkan bahawa kehadiran kelas-kelas HBP yang berbeza dalam

semen mungkin tidak boleh digunakan sebagai penunjuk kesuburan sesekor lembu jantan.

Heparin dan HBP yang dibeku kering didapati tidak mempunyai sebarang kesan ke atas peratus motiliti sperma lembu jantan Sahiwal-Friesian dan anab. Walaubagaimanapun, penambahan heparin atau HBP yang dibeku kering ke atas sperma epididimis dan sperma terejakulat telah memberikan penambahan yang signifikan dalam kadar reaksi akrosom kedua-dua spesis tersebut.

Kajian ini juga menunjukkan bahawa anab yang disuntik dengan sperma epididimis tanpa penambahan HBP menghasilkan min kadar kelahiran dan min kelahiran anak bagi seekor yang tinggi berbanding dengan ternakan-ternakan yang disuntik dengan sperma epididimis yang telah ditambah dengan kuantiti HBP yang berbeza. Keputusan-keputusan ini menunjukkan bahawa sperma epididimis anab boleh mempersenyawakan ovum tanpa didedahkan kepada HBP atau plasma semen terlebih dahulu, walaupun dipercayai bahawa HBP yang ada dalam plasma semen perlu untuk kapasitasi sperma lembu dan anab.

Kumpulan anab yang disuntik dengan sperma epididimis yang ditambah dengan HBP pada kadar 4 mg/ml, merekodkan min kelahiran anak bagi seekor yang rendah berbanding dengan anab betina yang disuntik dengan sperma epididimis yang ditambah dengan HBP pada kadar yang rendah. Keputusan-

keputusan ini mungkin menunjukkan bahawa kepekatan HBP yang tinggi membawa lebih keburukan kepada sperma epididimis daripada meningkatkan kesuburannya.

Di antara empat kumpulan arnab yang disuntik dengan sperma epididimis (ES), sperma epididimis ditambah dengan HBP (ESH), sperma terejakulat (ESM) dan spermatozoa terejakulat ditambah dengan HBP (ESMH), tiada perbezaan signifikan dalam kadar kelahiran anak antara kumpulan yang disuntik pada 0 jam selepas aruhan ovulasi. Walaubagaimanapun, bagi arnab-arnab yang disuntik pada 6 jam selepas aruhan ovulasi, kadar kelahiran adalah rendah yang signifikan dalam kumpulan yang disuntik dengan ES berbanding dengan ketiga-tiga kumpulan lain yang disuntik dengan sperma samada mempunyai HBP semulajadi atau HBP yang dibeku kering. Dalam kajian ini juga didapati bahawa tiada seekor pun arnab yang disuntik pada 12 dan 24 jam selepas aruhan ovulasi melahirkan anak.

Diperhatikan juga bahawa sampel-sampel sperma epididimis yang ditambah dengan HBP yang dibeku kering melekat lebih banyak heparin bertritium (^3H) pada membrannya berbanding dengan sampel-sample yang tidak ditambah dengan HBP. Kajian ini juga menunjukkan bahawa sperma dari lembu yang lebih subur melekat kepekatan heparin bertritium ^3H yang tinggi pada membran sperma berbanding dengan sperma dari lembu yang kurang subur.

Dirumuskan bahawa (i) sperma epididimis arnab boleh mempersenyawakan ovum tanpa didedahkan kepada HBP atau plasma semen terlebih dahulu pada arnab dan (ii) lembu jantan Sahiwal-Friesian yang lebih subur boleh dipilih berdasarkan keupayaannya melekat heparin bertritium ^3H pada membran sperma.

ACKNOWLEDGEMENTS

I am immensely grateful to Dato' Prof. Dr. Sheikh Omar Abdul Rahman for his guidance throughout my study at Universiti Putra Malaysia and for his invaluable comments and suggestions in the preparation of the thesis. His willingness to be my supervisor and his keen interest in my research projects are highly appreciated. My appreciation is also extended to Prof. M. R. Jainudeen and Dr. Nordin Yusof, co-supervisors, for their constructive ideas and suggestions during the preparation of the thesis.

I also would like to express my gratitude to the Director General, Malaysian Agricultural Research and Development Institute (MARDI) for allowing me to pursue the study at the university and providing the facilities to carry out the studies, and Director, Livestock Research Centre, for his support and encouragement throughout the course of my study.

Appreciation is also due to: Mr. Pullikutty Ayanar for his assistance in data collection and care of the experimental animals; En. Ahmad Shokri bin Othman for his help in the statistical analysis; and to my fellow research officers of the Livestock Research Centre, MARDI for their friendship, views and inspiration.

Last but not least, I would like to extend my sincere appreciation to my wife, Kathirvelu Mudaliar Pattu, children, brothers and sisters-in-law for their patience, understanding and sacrifice throughout the course of my study, and to my beloved parents and my family for their moral support.

I certify that an Examination Committee met on 30th December 2003 to conduct the final examination of Ramakrishnan a/l Patchaiappan on his Doctor of Philosophy thesis entitled "Influence of Heparin and Seminal Heparin Binding Proteins on Reproductive Performance of Sahiwal-Friesian Bulls and Rabbits" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Mohamed Ali Rajion, Ph.D.

Professor
School of Graduate Studies
Universiti Putra Malaysia
(Chairman)

Dato' Sheikh Omar Abdul Rahman, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

M.R. Jainudeen, Ph.D.

Professor Emeritus
Department of Veterinary Clinical Studies
Faculty of Veterinary Medicine
University Putra Malaysia
(Member)

Nordin Bin Yusof, Ph.D.

Research Officer
Stesen MARDI Bukit Ridan
Muadzam Shah
Pahang
(Member)

Ramli Yusof, Ph.D.

Professor
Institute of Biological Science
Faculty of Science
University of Malaya
Kuala Lumpur
(Independent Examiner)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 10 FEB 2004

The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Dato' Sheikh Omar Abdul Rahman, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

M.R. Jainudeen, Ph.D.

Professor Emeritus
Department of Veterinary Clinical Studies
Faculty of Veterinary Medicine
University Putra Malaysia
(Member)

Nordin Bin Yusof, Ph.D.

Research Officer
Stesen MARDI Bukit Ridan
Muadzam Shah
Pahang
(Member)

AINI IDERIS, Ph.D.
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

RAMAKRISHNAN A/L PATCHAIAPPAN

Date:

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	ix
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xviii
LIST OF FIGURES	xxi
LIST OF ABBREVIATIONS	xxii
CHAPTER	
1 INTRODUCTION	1
2 REVIEW OF LITERATURE	5
Male Reproduction	5
Molecular Mechanisms of Fertilization	5
Activation of Sperm Motility	6
Characteristics of Zona Pellucida	7
Spermatozoa Binding to the Zona Pellucida	8
Mechanism of the Acrosome Reaction	10
Fusion and Egg Activation	12
Libido and Mating Behavior in Bulls	14
The Assessment of Libido and Mating Behavior	16
Genetic	18
Breed Differences	18
Post Weaning Management	19
Hormone	20
Seasons and Enviromental Temperature	21
Frequency in Ejaculation	21
Nutrition	22
Environmental and Genetic Effects on Bull Fertility	22
Genetic Effects on Bull Fertility	23
Hormonal Influences	27
Seasons	27
Nutrition	29
Age	30



	Biochemical	31
	Assessment of Bull Fertility	31
	Breeding Soundness Examination (BSE)	32
	Libido and Sexual Behavior	35
	Heparin Binding Proteins from Seminal Plasma	37
3	ISOLATION, PURIFICATION AND CHARACTERISATION OF SEMINAL HEPARIN BINDING PROTEINS IN BULLS AND RABBITS	40
	Introduction	40
	Materials and Methods	42
	Animals and Management	42
	Vasectomy of Bulls	42
	Vasectomy of Rabbits	44
	Semen Collection and Seminal Plasma Separation in Bulls	44
	Semen Evaluation	45
	Progressive Motility	46
	Spermatozoa Concentration	46
	Semen Collection and Seminal Plasma Separation in Rabbits	46
	Grouping of Bulls According to their Fertility Status	47
	Isolation of Heparin Binding Proteins	48
	Characterisation of Heparin Binding Proteins	50
	Analysis of Data	51
	Results	52
	Discussion	61
4	MOTILITY AND ACROSOME REACTION OF RABBIT AND BOVINE SPERMATOZOA IN THE PRESENCE OF HEPARIN BINDING PROTEINS	66
	Introduction	66
	Materials and Methods	67
	Experiment 1	67
	Semen Collection, Evaluation and Extension of Bull Spermatozoa	68
	Selection	68
	Isolation of Epididymal Spermatozoa from Sahiwal-Friesian Bull	68
	In Vitro Induction of Acrosome Reaction	70
	Semen Extender	71
	Acrosomal Stain	71
	Staining Procedure	72

	Experiment 2	73
	Semen Collection and Evaluation	73
	Semen Evaluation	73
	Progressive Motility	74
	Spermatozoa Concentration	74
	Semen Extension	74
	Isolation and Extension of Rabbit Epididymal Spermatozoa	75
	In Vitro Induction of Acrosome Reaction in Rabbit Spermatozoa	76
	Analysis of Data	77
	Results	78
	Discussion	92
5	ARTIFICIAL INSEMINATION OF RABBIT USING EPIDIDYMAL SPERM AND EJACULATED SEMEN WITH AND WITHOUT THE ADDITION OF HEPARIN BINDING PROTEINS	97
	Introduction	97
	Materials and Methods	99
	Semen Collection, Seminal Plasma Separation and Isolation of Heparin Binding Proteins	99
	Isolation of Epididymal Spermatozoa	99
	Semen Extender	100
	Semen Collection and Evaluation	100
	Semen Evaluation	100
	Progressive Motility	101
	Spermatozoa Concentration	101
	Experiment 1	101
	To Determine the Influence of Four Levels IIBP On the Reproduction Performance of Rabbit	101
	Experiment 2	103
	A Comparative Study on the Effect of Epididymal and Ejaculated Spermtozoa Treated with IIBP on the Reproductive Performance of Rabbit.....	103
	Experiment 3	105
	To Evaluate Whether HBP Exert an Effect on Sperm to Alter their Fertilising Ability in Relation to Ovulation in Rabbit	105
	Analysis of Data	107

	Results	109
	Discussion	118
6	RELATIONSHIP BETWEEN BULL FERTILITY AND BINDING OF ³ H HEPARIN TO SPERMATOOZOA IN SAHIWAL-FRIESIAN BULL	125
	Introduction	125
	Materials and Methods	126
	Experiment 1	126
	Isolation of Epididymal Spermatozoa	126
	Semen Extender	127
	Semen Evaluation	127
	Progressive Motility	128
	Spermatozoa Concentration	128
	Heparin Binding Assay	128
	Experiment 2	132
	Grouping of Bulls According to their fertility Status	134
	Analysis of Data	134
	Results	136
	Experiment 1	136
	Experiment 2	147
	Discussion	150
7	GENERAL DISCUSSION	155
	Proposed Future Studies	163
8	SUMMARY AND CONCLUSION	165
	Summary	165
	Conclusion	167
9	BIBLIOGRAPHY	
10	LIST OF PUBLICATIONS FROM THIS THESIS	187
11	PHOTOGRAPHS OF SAHIWAL-FRIESIAN BULLS	189
12	BIODATA OF AUTHOR	190

LIST OF TABLES

Table	Page
2.1 Composite Breeding Soundness Examination (BSE) score	32
2.2 BSE values and pregnancy rates	34
2.3 Pregnancy rates and bull classification	35
2.4 Estimated genetic correlation between reproductive traits in bulls and age at puberty in half-sib heifers	36
3.1 Mean Concentration of Heparin Binding Proteins in Seminal Plasma of Bull	53
3.2 Mean Concentration of Heparin Binding Proteins in Seminal Plasma of Rabbit	55
3.3 Overall Mean Concentration of Heparin Binding Proteins in Seminal Plasma of Bull and Rabbit	57
3.4 Different classes of seminal heparin binding proteins in bull and rabbit	58
4.1 Motility of Epididymal Spermatozoa from Sahiwal-Friesian Bull Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	79
4.2 Motility of Ejaculated Spermatozoa from Sahiwal-Friesian Bull Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	80
4.3 Percentage of Induced Acrosome Reactions of Epididymal Spermatozoa from Sahiwal-Friesian Bull Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	82

4.4	Percentage of Induced Acrosome Reactions of Ejaculated Spermatozoa from Sahiwal-Friesian Bull Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	84
4.5	Viability of Rabbit Epididymal Spermatozoa Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	85
4.6	Viability of Rabbit Ejaculated Spermatozoa Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	87
4.7	Percentage of Induced Acrosome Reactions of Rabbit Epididymal Spermatozoa Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	89
4.8	Percentage of Induced Acrosome Reactions of Rabbit Ejaculated Spermatozoa Over a Period of 24 Hours in the Presence of Heparin and Lyophilised Heparin Binding Proteins	91
5.1	The Different Groups of Rabbits Inseminated with ES, ESH, ESM and ESMH at 0, 6, 12 and 24 Hours After Vaginal Stimulation	108
5.2	Effects of Four Levels of Heparin Binding Proteins on the Reproductive Performance of Rabbit.....	110
5.3	Effects of ES, ESH, ESM and ESMH on the Reproductive Performance in Rabbits	112
5.4	Effects of Epididymal and Ejaculated Spermatozoa on the Overall Mean (Irrespective of the HBP Treatment) Littered Rate, Litter Born Per Doe, and Gestation Period in Rabbit	114

5.5	Effects of Heparin Binding Proteins (Irrespective of Source of Spermatozoa) on the Overall Mean Littered Rate, Litter Born Per Doe, and Gestation Period in Rabbit	115
5.6	Effects of ES, ESH, ESM and ESMH on the Reproductive Performance in Rabbit Inseminated at 0 and 6 Hours After Vaginal Stimulation	116
6.1	Fertility status of nine Sahiwal-Friesian bulls	135
6.2	Mean Concentration of Tritiated Heparin (^3H) Bound to Epididymal Spermatozoa (Treated with and without Heparin Binding Proteins) of Sahiwal-Friesian Bulls (mean of 4 bulls) in a Twelve Serial Dilutions of Tritiated Heparin	144
6.3	Overall Mean* Concentration of Tritiated Heparin (^3H) Bound to Epididymal Spermatozoa (Treated with and without Heparin Binding Proteins) of Four Sahiwal-Friesian Bulls in a Twelve Serial Dilutions of Tritiated Heparin	145
6.4	Binding of ^3H heparin to spermatozoa of nine Sahiwal-Friesian bulls to a concentration of 1626 pmol of ^3H added	148
6.5	Tritiated Heparin Binding Abilities of Ejaculated Spermatozoa from Bulls of Low, Average and High Fertility using a ^3H heparin Concentration of 1626 pmol	149

LIST OF FIGURES

Figure		Page
3.1	The SDS-PAGE separation of heparin binding proteins of four Sahiwal-Friseian bulls of varying fertility	59
3.2	The SDS-PAGE separation of heparin binding proteins of vasectomised and unvasectomised rabbits	60
6.1	A representative saturation binding plot of (^3H) heparin bound to epididymal spermatozoa devoid of heparin binding protein in four Sahiwal-Friesian bulls	137
6.2	A representative saturation binding plot of (^3H) heparin bound to epididymal spermatozoa treated with heparin binding proteins in four Sahiwal-Friesian bulls.....	138
6.3	Asymptotic curves indicating the relationship between the predicted and overall mean (^3H) heparin bound to epididymal spermatozoa devoid of heparin binding proteins in Sahiwal-Friesian bulls	140
6.4	Asymptotic curves indicating the relationship between the predicted and overall mean (^3H) heparin bound to epididymal spermatozoa treated with heparin binding proteins in Sahiwal-Friesian bulls	142

LIST OF ABBREVIATIONS

AI	artificial insemination
AV	artificial vagina
BSE	breeding soundness examination
Ca	calcium
DNA	deoxyribonucleic acid
DPM	disintegration per minute
ES	epididymal spermatozoa
ESH	epididymal spermatozoa with heparin binding proteins
ESM	ejaculated spermatozoa
ESMH	ejaculated spermatozoa with heparin binding proteins
g	gram
GAG	glycosaminoglycan
GnRH	gonadotrophin releasing hormone
HBP	heparin binding proteins
³ H	tritiated heparin
hr	hour
IBHK	national animal biotechnology institute
IU	international unit
kDa	kilodaltons
kg	kilogram
M	molar concentration
MARDI	Malaysian agricultural research development institute
mCi	millicurie
min	minutes
ml	milliliter
mo	month
N	sample number
NRR	non-return rate
PBS	phosphate buffer saline
pH	logarithm of the reciprocal of the H ion concentration
rpm	rotation per minute
SDS	sodium dodecyl sulfate-polyacrylamide
SD	standard deviation
TALP	tyrode's albumin lactate pyruvate
TDN	total digestible nutrients
ZP	zona pellucida
μl	microliter



CHAPTER I

INTRODUCTION

The population of cattle and water buffalo in Peninsular Malaysia in 1998 was 742,171 and 148,786, respectively (Department of Veterinary Services, 2001). The industry is growing steadily over the years and has earned a place in national economy. The projected population increase for cattle by the year 2025 is estimated to be 993,639. This positive sign of growth in the cattle industry is expected to contribute more towards agriculture and food production.

Sperm capacitation and the acrosome reaction are requisite events in the process of fertilisation (Bedford, 1983). Heparin, one class of glycosaminoglycans (GAG) found in female reproductive tracts, was reported to induce capacitation and acrosome reactions in spermatozoa for bulls and rabbits (Lenz *et al.* 1983a, b). It has been reported that bovine ejaculated spermatozoa required nine hours of incubation with GAGs to undergo acrosomal reaction *in vitro*, whereas epididymal spermatozoa required 22 hours (Handrow *et al.* 1982; Lenz *et al.* 1982; Lee *et al.* 1985). However, a 20 minutes exposure of epididymal spermatozoa to seminal plasma was found to reduce the time required for the acrosome reaction to nine hours as the ejaculated spermatozoa (Lee *et al.* 1985). It was also found that if solubilised zonae pellucidae were used to induce acrosomal reaction in spermatozoa treated with heparin for four hours, ejaculated

spermatozoa responded with acrosome reaction but epididymal spermatozoa did not (Florman and First, 1988a; Florman *et al.* 1989). The exposure of epididymal spermatozoa to seminal plasma *in vitro* enable those spermatozoa to be capacitated by heparin and to respond to zonae pellucidae with an increase in acrosome reaction in a manner similar to ejaculated spermatozoa (Florman and First, 1988). Lee et al. (1985) further showed that seminal plasma increased the number of binding sites for heparin on epididymal spermatozoa. Miller *et al.* (1987) found that bovine seminal plasma which contains several proteins known as heparin binding proteins (HBP), upon ejaculation bound to spermatozoa and were responsible for the heparin bind ability of the spermatozoa. There is scanty information on the effect of purified seminal HBP on the ejaculated and epididymal spermatozoa in Sahiwal-Friesian bull and rabbit. There is also a lack of information on whether purified seminal HBP exert an effect on spermatozoa to alter their fertilising ability in relation to ovulation in rabbit.

Although it has been shown that seminal plasma had increased the number of heparin binding sites on epididymal spermatozoa (Lee et al. 1985), no study has been carried out to determine whether the purified HBP could bring about the same reaction in epididymal spermatozoa. Furthermore, there is lack of information on the amount of heparin bound to the epididymal and ejaculated spermatozoa treated with purified seminal HBP.